


Cystamine dihydrochloride



CAS
56-17-7

Status
Commercial

Inquiry

Cystamine dihydrochloride belongs to the group of aminothiols. The active substance binds the cysteine in the lysosomes. It is used for the treatment of cystinosis, cystinuria and radiation sickness.

Description

Formula C₄H₁₄Cl₂N₂S₂

Molecular weight 225.20 g/mol

Molecular size small

Controlled Substance no

Precursor no

Your Contact



Marius Hang
International Division
Director API Export

Midas Pharma GmbH
Rheinstr. 49
55218 Ingelheim
Germany

Disclaimer

The information contained herein is the proprietary information of Midas Pharma GmbH and its affiliates. This information is provided to you solely for your evaluation, to determine your interest in a potential business relationship with Midas Pharma GmbH and/or its affiliates. Midas Pharma GmbH excludes any express or implied representations, guarantees or warranties of whatsoever nature, including but not limited to the warranties for the completeness or accuracy of the information provided herein.

No information provided herein – including any reference to any product or service – constitutes an offer for sale, or shall be construed as representing an offer for sale. In particular, nothing herein should be construed as a promotion or advertisement for any product, or for the use of any product, that infringes valid patents and/or is not authorized by the laws and regulations of your country of residence.